

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266212376>

The lichen flora of the Chagos Archipelago, including a comparison with other island and coastal tropical floras

Article · December 2000

DOI: 10.11646/bde.18.1.22

CITATIONS

16

READS

16

2 authors:



Mark Seaward

University of Bradford

175 PUBLICATIONS 2,792 CITATIONS

[SEE PROFILE](#)



Andre Aptroot

Adviesbureau voor Bryologie en Lichenologie

470 PUBLICATIONS 7,985 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Lichen Flora of Iran, An International Project [View project](#)



Notes for genera in Ascomycota [View project](#)

The lichen flora of the Chagos Archipelago, including a comparison with other island and coastal tropical floras

Mark R.D. Seaward

Department of Environmental Science, University of Bradford, Bradford BD7 1DP, UK

André Aptroot

Centraalbureau voor Schimmelcultures, P.O.Box 273, 3740 AG Baarn, The Netherlands

Abstract. The 1996 Chagos Expedition provided the first opportunity to study the archipelago's lichen flora. Seventeen of the 55 islands were ecologically investigated, some in more detail than others, and lists and representative collections of lichens have been assembled for many of them. In all, 67 taxa have been recorded, 52 to specific level.

Although the islands have a low biodiversity for cryptogamic plants, as would be expected in terms of their relatively young age, remoteness and small terrestrial surface areas, those taxa that are present are often found in abundance and play significant ecological roles.

There is a good correlation between total lichen biodiversity and island size, despite the fact that *Cocos nucifera* is such an important substratum for cryptogamic plants and its presence on all islands studied provides a consistently high associated species count.

Comparisons of lichen floras for ten island and coastal tropical areas show good correlations (based on the Sørensen Coefficient) within the Indian Ocean as would be expected, but poorer correlations exist within and between Pacific Ocean and neotropical floras. Ranked correlations between Chagos and other floras are in the sequence Maldives > Laing Island > Aldabra > Tuamotu > Pitcairn > N.Mariana & Belize > Guadeloupe > Cook. When coefficients are calculated using only the Physciaceae, different correlations and sequences are derived, but the affinities of the Indian Ocean islands remain strong. However, although the lichen flora of Chagos is characteristic for an Indian Ocean, it is dominated by pantropical species.

Introduction

The Chagos Archipelago is the largest and most isolated atoll complex in the world. Although the actual archipelago covers c.37500 km² of mainly coral reefs, only about 2% provides terrestrial habitats. The 55 islands of the archipelago (Figure 1) have a typical tropical maritime climate, with an annual rainfall of about 2400 mm and an average temperature of c.27°C. Recent research (Eisenhauer *et al.*, 1999) has shown that the current islands vary in age between 2800 and 6500 years. For further information on the history, reef formation, climate, fauna and flora of these islands see Sheppard and Seaward (1999).

The islands were exploited for copra from the 1790s onwards; much of the native vegetation was removed to make way for *Cocos* plantations and many of the islands were inhabited by plantation workers. The plantations were run down in 1967 as their commercial future could not be guaranteed, and by 1973 the islands had become uninhabited, apart from the largest, Diego Garcia which supports a military base.

Diego Garcia (44 km²) is a horseshoe-shaped atoll enclosing a large, deep lagoon with three small islands at its mouth. It was made available for British and USA defence purposes in 1966 and normally supports a military base of c.3400 personnel. The other main island groups of the archipelago are the Peros Banhos atoll (29 islands, total land area of 10.4 km²) and the Salomon atoll (11 islands, total land area of 5.2 km²). The Great Chagos bank comprises eight islands widely separated from the other atolls. In addition, the Egmont atoll, formerly composed of six islands but now, as observed by the 1966 expedition members, evidently coalescing to form only three, is the smallest of the island groups.

Since their abandonment, most islands have been rarely visited, but extended visits by yachtsmen to the Salomon and Peros Banhos atolls are becoming increasingly common. The few other visitors have been mainly scientists, the most recent being the 1996 Chagos Expedition, permission for which centred on the requirement by the authorities for a conservation management plan, with much of the emphasis placed on an assessment of the status of the archipelago in biogeographical terms. Much of the research

concentrated on the coral reefs and marine fauna, but two entomologists (Linda Barnett and Craig Emms) and three botanists (David Bellamy, Ursula Gregory and Mark Seaward) worked on terrestrial systems. Preliminary botanical work by Ursula Gregory on several islands of the Peros Banhos and Salomon atolls during early stages of the expedition was extended by more detailed studies, employing taxonomic, ecological and remote sensing techniques, made by the two other botanists who were fortunate in being able to visit a high proportion of the archipelago's islands.

Ecology

Despite their relatively young age and small terrestrial surface area, the islands have a relatively high biodiversity in terms of higher plants. Currently there are c.280 species present, but only c.35 of these comprise the original native flora prior to human arrival (Topp & Sheppard, 1999). Since the cryptogamic plants had not previously been given serious attention, there being only a short published list of mosses (Townsend 1971), Mark Seaward (with supplementary work by Ursula Gregory and Fred Stewart) surveyed the lichen and bryophyte floras of many of the islands, and also made collections of cyanobacteria, algae and fungi. Although there is a relatively low diversity for most of these groups in terrestrial habitats, and any 'new' species discovered there will undoubtedly occur elsewhere, they are abundant and ecologically important.

Due to their small size and difficulty in identification, particularly *in situ*, more remain to be discovered, but to date only 9 moss, 4 liverwort, 68 lichen, 1 algal and 26 cyanobacterial taxa have been determined from strictly terrestrial habitats; a rich fungal flora undoubtedly exists, but due to the limited material available for study, only 21 macrofungal, 5 microfungal and 1 myxomycete taxa have been identified so far (Seaward 1999).

Lichens on Chagos would appear not to grow on coral, or on the poorly developed sandy soils arising from it, with the exception of an unnamed *Siphula* sp. which colonizes several square metres of exposed soil (pH c.7.4) on a single island. However, cyanobacteria, which extensively clothe eroded corals and sandy soils,




Fig. 1. Chagos Archipelago (see Table 1 for details of islands lichenologically studied).

provide a continuously renewable nitrogen and carbon source and improve soil stability through particle aggregation. With time, such soils may support lichens, but in the meantime their role is upheld by the cyanobacterium *Nostoc commune* which covers very extensive areas of exposed soils, its presence only fully appreciated after rain.

However, bryophytes are important in stabilising soils and appear to be the primary stages in the succession to higher vegetation in exposed areas of the islands. The soil around and below mosses is colonized by cyanobacteria which release not only substantial amounts of fixed nitrogen but also growth substances which benefit the moss (Rogers & Henriksson 1976, Richardson 1981). The reduction or loss of breeding colonies of birds from islands due to rat infestation will lead to reduced inputs of phosphorus and nitrogen which will ultimately affect the capacity of some cryptogamic plants to establish themselves on soils inherently poor in nutrients.

Man-made building materials are similarly devoid of lichens, but cyanobacteria are to be found in abundance and relatively high diversity, their presence creating a characteristic black coating over the stonework, etc. of abandoned plantation buildings. *Scytonema-Gloeocapsetum*, the commonest association encountered in the Chagos islands, is well developed on such substrata where exposed to sunlight and wetted sporadically (Hoffmann 1989).

A rich cyanobacterial flora on the bark of many trees is supported by one or more as yet undetermined species of the alga *Trentepohlia*. Of the epiphytic floras studied to date, it would appear that *Casuarina equisetifolia* supports the largest diversity of cyanobacteria; *Cocos nucifera*, on the other hand, has only a few cyanobacteria species.

However, lichens readily colonize tree bark surfaces, particularly in the case of *Cocos nucifera*, its smooth, hard and usually dry bark supporting a rich lichen flora often dominating large areas of the trunk; however, there is a low diversity and poor coverage of bryophytes, except about the tree bases, particularly those subjected to higher humidity levels where cushions of the moss *Calymperes tenerum* can be found in plenty.

As Sipman and Harris (1989) note, lichens in tropical rain forests show little preference among phorophytes: this is amply demonstrated by the 20 or so phorophyte species studied on Chagos, with the exception of *Cocos nucifera* which due to its frequency appears to have developed its own distinctive epiphytic flora.

A lichen zonation from base to crown (cf. Sipman & Harris 1989) is not distinguishable on Chagos trees, and although there are some differences between the epiphytic floras of the base, trunk and crown, zones are often delimited vertically according to aspect. In habitats frequented by birds, a natural lichen mosaic is often ill-defined due to the presence of other lichen species which are encouraged by the nutrient enrichment of tree trunks and branches. A 'nutrient rain' rich in leachates of excrement (Sipman & Harris 1989) derived from tree canopy birds, their status and behaviour governed by the presence/absence of rats on particular islands, greatly influences the lichen flora.

The nature of the epiphytic flora changes with age, and as the trunks of all species die, different bryophytes support the decomposition process, the rotting wood community consisting mainly of *Ectropothecium* species (cf. Pocs 1982; Seaward 1999). Although a diversity of colourful fungi were also present, often forming a feature of the characteristic crater-shaped tree bases left when coconut palms topple over, lichens appeared not to be associated with this muscicolous habitat.

Two foliicolous lichens, *Opegrapha vegae* and *Porina perminuta*, are commonly found on *Cocos* leaves, which are also colonized by microfungi, such as *Mycosphaerella gastonis*; these become much more prevalent with time and a new microflora takes over once the leaves have been shed. The presence of foliicolous lichens on leaves other than those of *Cocos* and *Calophyllum inophyllum* was not investigated.

Fieldwork

Those terrestrial locations visited on the 1996 Expedition are shown in Figure 1 and listed in Table 1. In all, 25 islands were visited, but only 17 were investigated and collections, albeit variable

Table 1. Islands of the Chagos Archipelago lichenologically studied during the 1996 Expedition

Code	Location	Size (ha)	Lichen Taxa*	Collectors**
B	Three Brothers:			
B(N)	North	6.0	11	[1]
B(M)	Middle	8.0	15	[1]
B(S)	South	23.0	17	[1]
DG	Diego Garcia	2719.5	43	[1]
DG(S)	Diego Garcia (settlement)			[1]
DG(I)	Diego Garcia (East, Middle & West Islands)	14.5		[3]
E	Eagle Island(s)	243.5	20	[1]
EG	Egmont:			
EG(L)	Lubine	177.0	18	[1]
EG(S)	Sipaille	64.0	15	[1]
N	Nelson Island	81.0	13	[1]
PB	Peros Banhos:			
	Ile du Coin	127.5		[2]
	Ile Diamante	88.5		[2]
	Ile Yeye	58.5		[2]
S	Salomon Islands:			
SI(A)	Ile Anglaise	75.5	15	[1,2]
SI(B)	Ile Boddam	108.5	32	[1,2]
SI(D)	Ile Diable	0.1	8	[1]
SI(DP)	Ile de la Passe	26.0	10	[1,2]
SI(DS)	Ile du Sel	2.0		[2]
SI(F)	Ile Fouquet	39.5	21	[1,2]
SI(J)	Ile Jacobin	1.5	8	[1,2]
SI(M)	Ile Mapou	4.0	10	[1,2]
SI(P)	Ile Poule	0.5		[2]
SI(S)	Ile Sepulture	0.5	14	[1]
SI(T)	Ile Takamaka	44.0	15	[1,2]

* Minimal no. of taxa determined.

** [1] M.R.D.Seaward, [2] Ursula Gregory and [3] F.Stewart.

in size and diversity, made of different groups of cryptogamic plants. Collections from East, Middle and West Islands of Diego Garcia, all islands of Peros Banhos, and Ile Poule and Ile du Sel of the Salomon Islands are limited in scope, the author having been unable to visit these (as well as Danger Island, Sea Cow and Resurgent) in order to make field notes and collect representative material.

Based on the fieldwork undertaken during

the 1996 Expedition, the lichen flora list which follows is almost complete, and an attempt has been made to determine the status of the taxa for each of the islands and their relationship to (1) neighbouring islands within the archipelago, (2) other islands in the Indian Ocean, and (3) to other island and coastal tropical floras of the world. Relationships between the Chagos lichen flora and those of the mainlands of Africa, India and South-east Asia have yet to be determined (cf.

Table 2. Relationships between ten island and coastal lichen floras based on Sorensen coefficients (K) for (a) all taxa and (b) only Physciaceae. Sources: Aldabra - Seaward *et al.* (1996); Belize - Barclay-Estrup (1992); Chagos - Seaward (1999); Guadeloupe - Aptroot (1998, unpublished list); Laing Island - Aptroot *et al.* (1995); Maldives - Aptroot (1991); Cook, N.Mariana, Pitcairn & Tuamotu - Elix & McCarthy (1998).

Contrasting Locations	(a)		(b)
	K	Rank	Rank
<i>Good correlation:</i>			
Maldives - Aldabra	34.5	1	1
Chagos - Maldives	32.8	2	3
<i>Reasonable correlation:</i>			
Tuamotu - Pitcairn	24.5	3	3
Chagos - Laing Is	24.1	4	3
N.Mariana - Cook	23.3	5	2 5
Chagos - Aldabra	22.6	6	9
Aldabra - Tuamotu	22.2	7	1 0
N.Mariana - Pitcairn	22.0	8	2
Chagos - Tuamotu	21.6	9	3
N.Mariana - Tuamotu	19.8	10	3
Maldives - Tuamotu	19.5	11	1 5
Chagos - Pitcairn	18.7	12	7
Maldives - Laing Is	18.7	12	2 9
<i>Moderate correlation:</i>			
Maldives - Guadeloupe	15.1	14	2 2
Cook - Pitcairn	14.8	15	2 7
Maldives - Pitcairn	14.3	16	1 6
Maldives - Belize	14.3	16	3 7 *
Chagos - N.Mariana	13.1	18	1 1
Chagos - Belize	13.1	18	3 5
Maldives - Cook	13.0	20	1 1
Aldabra - Guadeloupe	12.5	21	1 1
Chagos - Guadeloupe	11.6	22	3 5
Tuamotu - Belize	11.4	23	2 9
Pitcairn - Belize	11.1	24	1 6
Aldabra - Pitcairn	10.8	25	2 1
Laing Is - Guadeloupe	10.6	26	3 7 *
<i>Weak correlation:</i>			
Aldabra - Cook	9.8	27	1 8
Laing Is - Tuamotu	9.8	27	1 4
Aldabra - N.Mariana	8.7	29	1 8
Laing Is - N.Mariana	8.7	29	2 2
Aldabra - Belize	8.7	29	2 7
Aldabra - Laing Is	8.6	32	1 8
Guadeloupe - Belize	8.5	33	3 7 *
Chagos - Cook	7.6	34	3 4
Tuamotu - Cook	7.5	35	2 5
Laing Is - Belize	5.8	36	2 9
Maldives - N.Mariana	5.4	37	2 9

Poor correlation:

Laing Is	Pitcairn	4.8	38	24
Cook	Guadeloupe	3.1	39	37*
N.Mariana	Belize	2.9	40	29
Laing Is	Cook	2.3	41	37*
N.Mariana	Guadeloupe	2.2	42	37*

No correlation:

Tuamotu	Guadeloupe	0	43	37*
Pitcairn	Guadeloupe	0	43	37*
Cook	Belize	0	43	37*

* no correlation

Renvoize 1975).

The relationship between biodiversity and island size has been investigated: preliminary studies show that there is a good correlation ($r = 0.7465$) between the total number of lichen taxa and island size (Figure 2), despite the fact that *Cocos* is such an important substratum for lichens and its presence on all the islands studied provides a consistently high associated species count.

As a result of the 1996 Expedition, it was hoped to establish credible baseline data for future studies aimed at monitoring lichen stability in the light not only of local disturbance but also far-ranging global impacts. Although this may be possible for the islands' terrestrial habitats, it should be noted that lichens did not occur on exposed coral of the island fringes: zonal patterns capable of providing accurate monitors of seawater pollution and of rising sea-levels were lacking. The algal and cyanobacterial floras of the marine and tidal areas of the islands, which are undoubtedly much more biodiverse and contain many native, and indeed endemic, species, may be better suited to monitor such changes.

List of Taxa Recorded

Sixty-seven taxa were recorded, 52 and 15 determined to specific and generic level. The taxonomic list provides in the following sequence (1) some indication of frequency of occurrence; (2) preferred substrata; (3) distribution in the

Chagos Archipelago (according to the abbreviations given under 'Locations Investigated' above); and (4) world distribution where known. Specimens of all the taxa recorded by Mark Seaward are in Herb.Seaward, and in the case of collections made by Ursula Gregory in Herb.E; duplicates of some material collected by the author have been deposited in Herb. Aptroot, B, BG, BM, E, Ekman, H, IMI, Lücking, MO and OXF.

Agonimia papillata (O.Eriksson) Diederich & Aptroot Occasional, on *Hernandia sonora* & bases of *Pisonia grandis*.

B(M); B(S). Pantropical.

Herb. Aptroot, OXF, MRDS 107322, 107506

Anisomeridium subprostans (Nyl.) R.C.Harris

Rare, on *Hernandia sonora*.

SI(F). Pantropical.

Herb. MRDS 107418

Arthonia antillarum (Fée) Nyl.

Locally frequent on *Scaevola sericea*; less frequent on *Argusia argentea*, *Cocos nucifera*, *Guetarda speciosa* & *Hernandia sonora*.

B(M); B(N); B(S); DG; DG(I); EG(L); EG(S); N; SI(DP). Pantropical.

Herb. Aptroot, MRDS 107371, 107399, 107430, 107434

Arthonia sp.A

[cf. *A.ravida* Stirton]

Rare, on *Cocos nucifera*.
B(S).
Herb. MRDS 108367

Arthonia sp.B

on *Casuarina equisetifolia* & *Cocos nucifera*.

DG; EG(L).
Herb. IMI 374554

Arthopyrenia majuscula (Nyl.) Zahlbr.

Locally frequent, particularly on *Scaevola sericea*; less frequent on *Barringtonia speciosa*, *Casuarina equisetifolia*, *Cocos nucifera*, *Guettarda speciosa* & *Premna obtusifolia*.

B(M); B(N); B(S); DG; E; EG(L); EG(S); N; SI(A); SI(B); SI(D); SI(F). Pantropical.

Herb. Aptroot, E, OXF, MRDS 107325, 107326, 107354, 107355, 107398, 107427, 107508, 107513

Bacidia hostheleoides (Nyl.) Zahlbr.

Frequent, on *Argusia argentea*, *Cocos nucifera*, *Guettarda speciosa*, *Hernandia sonora*, *Scaevola sericea* & lignum.

B(M); DG; DG(I); E; EG(L); EG(S); N; SI(A); SI(B); SI(F); SI(J); SI(S); SI(T). Neotropical.

Herb. BM, MRDS 108120, 108121, 108122, 108805

Bacidia medialis (Nyl.) Zahlbr.

Rare, on *Argusia argentea* & lignum.
DG, DG(I), N, SI(J). Pantropical.

Herb. MRDS 108123, 108800, 108880

Bacidia sp.

On *Argusia argentea*, *Cocos nucifera* & *Hernandia sonora*.

B(N); DG; DG(I); EG(L); N; SI(S); SI(T).

Herb. MRDS 108801, 108802

Buellia efflorescens Müll.Arg.

Common on *Argusia argentea*, *Cocos nucifera* & *Morinda citrifolia*.

B(M); B(N); B(S); DG; DG(I); E; EG(L); EG(S); N; SI(A); SI(B); SI(F); SI(S). Pantropical.

Herb. MRDS 107352, 107353, 107364, 107375, 107502, 107575, 107581

Byssolecania fumosonigricans (Müll.Arg.) R.Sant.

Rare, on green stem of unidentified plant.
DG. Pantropical.

Herb. MRDS 108881

Calicium hyperelloides Nyl.

Rare, on *Cocos nucifera*.

SI(B). Pantropical to subtropical.

Herb. MRDS 107323

Caloplaca sp.

Rare, on *Casuarina equisetifolia* & *Scaevola sericea*.

DG; EG(L); EG(S).

Herb. MRDS 108803

Chrysothrix candelaris (L.) Laundon

Uncommon, on *Casuarina equisetifolia*.

DG; SI(B). Cosmopolitan.

Herb. Aptroot, B, BG, E, OXF, MRDS 107340, 107392, 107540, 107545

Coccocarpia cf. domingensis Vainio

Rare, on *Cocos nucifera*.

SI(B). Pantropical.

Herb. MRDS 107351

Coccocarpia erythroxyli (Sprengel) Swinscow & Krog

Rare, on *Cocos nucifera*.

SI(B). Pantropical to subtropical.

Herb. MRDS 107383

Coccocarpia palmicola (Sprengel) Arvidsson & Galloway

Locally frequent on *Cocos nucifera*, *Casuarina equisetifolia* & *Terminalia catappa*.

DG; E; EG(L); SI(B); SI(DS). Pantropical.

Herb. Aptroot, B, BM, E, OXF, MRDS 107321, 107324, 107339, 107391, 107421, 107515, 107576, 108068

Coccocarpia rottleri (Ach.) Arvidsson

Rare, on *Cocos nucifera*.

SI(B). Pantropical.

Herb. MRDS 107621

Coenogonium sp.

Locally frequent on stumps, adventitious

roots & husks of *Cocos nucifera*; apothecia infrequent.

B(M); DG; E; N; SI(B); SI(J).

Collema rugosum Krempelh.

Locally frequent on *Artrocarpus atilis*, *Barringtonia speciosa*, *Calophyllum inophyllum*, *Casuarina equisetifolia*, *Cocos nucifera*, *Cordia subcordata*, *Hernandia sonora*, *Hibiscus tiliaceus*, *Intsia bijuga*, *Morinda citrifolia*, *Premna obtusifolia* & *Terminalia catappa*.

DG; DG(S); SI(A); SI(B); SI(D); SI(DS); SI(F); SI(S); SI(T) - probably on all islands. Palaeotropical.

Herb. MRDS 107318, 107319, 107348, 107373, 108109

Cryptolechia subincolorella (Nyl.) D.Hawksw.

Infrequent, on *Hernandia sonora*, *Intsia bijuga*, *Morinda citrifolia*, *Pisonia grandis* & *Tabebuia pallida*.

B(S); DG; E; SI(B); SI(F); SI(S); SI(T). Paleotropical.

Herb. OXF, MRDS 107317, 107349, 107350, 107388

Dimerella sp.

Rare, on adventitious roots of *Cocos nucifera* & lignum.

DG; SI(J).

Herb. MRDS 108804

Dirinaria applanata (Fée) Awasthi

Infrequent, on *Barringtonia speciosa*, *Casuarina equisetifolia* & *Cocos nucifera*.

DG; SI(F); SI(S). Pantropical.

Herb. MRDS 107374, 107514

Dirinaria picta (Sw.) Clem. & Shear

Common, on *Artrocarpus atilis*, *Barringtonia speciosa*, *Calophyllum inophyllum*, *Casuarina equisetifolia*, *Cocos nucifera*, *Intsia bijuga* & *Scaevola sericea*.

B(S); DG; E; EG(L); EG(S); SI(A); SI(B); SI(D); SI(DP); SI(F); SI(M); SI(S); SI(T) - probably on all islands. Pantropical.

Herb. Aptroot, E, OXF, MRDS 107330, 107333, 107335, 107336, 107338, 107342, 107393, 107405, 107420, 107423, 107424,

107500, 107505, 107559

Endocarpon pusillum var. pallidum (Ach.) Körber

Rare, on cementwork of wall-tops, abandoned plantation buildings and stone memorials.

DG; SI(B). Cosmopolitan.

Herb. MRDS 107312, 107313

Fellhanera subtilis (Vezda) Diederich & Sérusiaux

Rare, on ? *Cocos nucifera*.

DG; DG(I).

Widespread (mainly European, but probably cosmopolitan).

Herb. MRDS 107737

Graphina analoga (Nyl.) Zahlbr.

Uncommon, on *Cocos nucifera* & *Terminalia catappa*.

SI(B); SI(F). Pantropical.

Herb. MRDS 107345, 107346

Graphina streblocarpa (Bel.) Müll.Arg.

Infrequent, on *Cocos nucifera* (particularly about base on adventitious roots) & *Melifera indica*.

DG; SI(B). Pantropical.

Herb. OXF, MRDS 107344, 107347

Graphis lineola Ach.

Infrequent, on *Barringtonia speciosa* & *Casuarina equisetifolia*.

DG; SI(F). Pantropical.

Herb. OXF, MRDS 107554, 107561, 108366

Graphis sp(p).

on *Barringtonia speciosa*(?) & *Casuarina equisetifolia*.

DG; DG(S); EG(S); SI(B); SI(DP).

Hyperphyscia syncolla (Nyl.) Kalb

Rare, on *Cocos nucifera*.

B(M). Pantropical.

Herb. MRDS 107435

Lecanora achroa Nyl.

Occasional on *Barringtonia speciosa*, *Casuarina equisetifolia* & *Cocos nucifera*.

DG; SI(B); SI(F). Pantropical.

Herb. E, OXF, MRDS 107794, 107795, 107796

Lecanora ecoronata Vainio

Occasional on *Guettarda speciosa* & *Scaevola sericea*.

B(S); E; EG(L); EG(S); SI(B); SI(F). Probably pantropical.

Herb. MRDS 107792, 107793

Lecanora sp.

[cf. *L.expallens* Ach.]

Frequent, on *Argusia argentea*, *Casuarina equisetifolia*, *Cocos nucifera* & *Scaevola sericea*.

DG; DG(I); EG(L); EG(S); SI(A); SI(B); SI(DS); SI(T).

Herb. E, MRDS 107881, 107882, 107885

Lecanora leproplaca Zahlbr.

Frequent, on *Cocos nucifera*, *Cordia subcordata* & *Premnea obtusifolia*

DG; E; EG(L); SI(A); SI(B); SI(J). Pantropical.

Herb. MRDS 107797, 107798

Lecanora tropica Zahlbr.

Rare, on *Cocos nucifera*.

E. Pantropical.

Herb. MRDS 107791

Lecanora sp.

Perhaps an undescribed sp. [cf. *L.usambarensis* Müll.Arg.]

Rare, on *Casuarina equisetifolia*

DG.

Herb. MRDS 107883

Lithothelium hyalosporum (Nyl.) Aptroot

Uncommon, on *Cordia subcordata* & *Hernandia sonora*.

SI(F); SI(M). Cosmopolitan.

Herb. MRDS 107368, 107417

Lithothelium illotum (Nyl.) Aptroot

Rare, on *Intsia bijuga*.

SI(T). Pantropical.

Herb. MRDS 107620

Nadvornikia hawaiiense (Tuck.) Tibell

Rare, on *Cocos nucifera*.

DG. Pantropical.

Herb. Aptroot, OXF, MRDS 107377, 107378

Opegrapha vegae R.Sant.

Common, on *Cocos nucifera* leaves.

B(M); B(N); B(S); DG; EG(L); EG(S); N; SI(A); SI(B); SI(DP); SI(F); SI(S); SI(T) - probably on all islands. Pantropical.

Herb. Aptroot, B, BG, BM, E, Lücking, OXF, MRDS 107307, 107308, 107309, 107310, 107311, 107412, 107413, 107414, 107415

Opegrapha sp.A

[cf. *O.confertoides* Torrente & Egea]

Rare, on *Cocos nucifera*.

SI(DP).

Herb. MRDS 108341

Opegrapha sp.B

[cf. *O.rufescens* Pers.]

Uncommon, on *Argusia argentea*, *Cocos nucifera* & *Herandiasonora*.

B(M); B(N); B(S); N.

Herb. MRDS 108338, 108340, 108342, 108345, 108355

Opegrapha sp.C

[cf. *O.viridis* (Ach.) Nyl.]

Common, on *Argusia argentea*, *Calophyllum inophyllum*, *Casuarina equisetifolia*, *Cocos nucifera*, *Cordia subcordata*, *Hernandia sonora*, *Hibiscus tiliaceus*, *Intsia bijuga*, *Morinda citrifolia*, *Pisonia grandis* & *Tabebuia pallida*; on all islands studied.

Herb. Aptroot, E, MRDS 108339, 108343, 108344, 108346, 108349, 108350, 108351, 108352, 108356, 108357, 108359

Opegrapha sp.D

[cf. *O.vulgata* (Ach.) Ach.]

Rare, on *Cocos nucifera* & *Scaevola sericea*.

SI(D); SI(DP).

Herb. MRDS 108347, 108358

Opegrapha sp.E

Infrequent, on *Argusia argentea*, *Calophyllum inophyllum*, *Casuarina equi-*

setifolia, *Cocos nucifera*, *Cordia subcordata*, *Hernandia sonora*, *Hibiscus tiliaceus* & *Intsia bijuga*.

B(M); DG; DG(I); E; EG(L); EG(S); SI(A); SI(B); SI(F); SI(M); SI(T).

Herb. MRDS 108347, 108358

Phaeographis sericea (Eschw.) Müll.Arg.

Uncommon, on *Casuarina equisetifolia*.
DG. Pantropical.

Herb. MRDS 108799

Physcia cf. dimidiata (Arnold) Nyl.

Rare, substratum unidentified.
SI(D). Cosmopolitan.

Physcia erumpens Moberg

Uncommon, on *Cocos nucifera* & *Hernandia sonora*, often over a cushion of moss (*Calymperes* spp.).

DG - ? restricted to one island. Pantropical.
Herb. MRDS 107342, 107343

Physcia integrata Nyl.

Locally frequent on *Cocos nucifera*; less common on *Guettarda speciosa* & *Hernandia sonora*.

B(M); B(N); B(S) - ? restricted to The Three Brothers. Pantropical.

Herb. Aptroot, OXF, MRDS 107384, 107403, 107432, 107517

Physcia krogiae Moberg

Rare, on *Pisonia grandis*.

B(S). Widespread in tropical & subtropical areas.

Herb. OXF, MRDS 107358

Physcia solediosa (Vainio) Lynge

Common on *Calophyllum inophyllum*, *Casuarina equisetifolia*, *Cocos nucifera*, *Cordia subcordata*, *Guettarda speciosa*, *Hernandia sonora*, *Intsia bijuga* & *Scaevola sericea*; uncommon on cement & rare on ironwork.

DG; E; EG(L); EG(S); SI(A); SI(B); SI(D); SI(DP); SI(DS); SI(F); SI(J); SI(M); SI(S); SI(T) - probably on all islands. Pantropical.

Herb. Aptroot, B, BM, E, H, OXF, MRDS 107327, 107328, 107334, 107341, 107356, 107357,

107367, 107370, 107389, 107394, 107395, 107396, 107397, 107406, 107416, 107419, 107499, 107501, 107503, 107560

Porina perminuta Vainio

Common on *Cocos nucifera* leaves; locally frequent on *Calophyllum inophyllum* leaves.

B(M); B(N); DG; E; EG(L); EG(S); N; SI(A); SI(B); SI(F); SI(J); SI(S) - probably on all islands. Mainly Pacific islands.

Herb. Aptroot, B, BG, BM, E, H, Lücking, OXF, MRDS 107301, 107302, 107303, 107304, 107305, 107306, 107407, 107408, 107409, 107410, 107411, 107757

Pyrenula concatervans (Nyl.) R.C.Harris

Locally frequent on *Calophyllum inophyllum*, *Guettarda speciosa*, *Hibiscus tiliaceus*, *Pisonia grandis* & *Tabebuia pallida*; frequent on *Intsia bijuga*.

B(S); DG; E; SI(A); SI(B); SI(DS); SI(F); SI(S); SI(T). Pantropical.

Herb. Aptroot, E, MRDS 107362, 107365, 107366, 107369, 107385, 107558

Pyrenula confinis (Nyl.) R.C.Harris

Uncommon, on *Casuarina equisetifolia* & *Premna obtusifolia*.

DG. Pantropical.

Herb. MRDS 107390, 107504, 107516, 107556

Pyrenula macularis (Zahlbr.) R.C.Harris

Uncommon, on *Casuarina equisetifolia* & *Intsia bijuga*.

DG; SI(T). Pantropical.

Herb. MRDS 107361, 107510

Pyrenula nitidula (Bres.) R.C.Harris

Rare, on *Hernandia sonora*.

DG. Pantropical.

Herb. OXF, MRDS 107431

Pyrenula ochraceoflava (Nyl.) R.C.Harris

Abundant on *Cocos nucifera*; less common on *Barringtonia speciosa*, *Calophyllum inophyllum*, *Casuarina equisetifolia*, *Hernandia catappa*, *Pisonia grandis*, *Premna obtusifolia* & *Scaevola sericea*.

B(M); B(N); B(S); DG; E; EG(L); EG(S); N; SI(A); SI(B); SI(D); SI(DP); SI(F); SI(M); SI(S); SI(T). On all islands studied. Pantropical.

Herb. Aptroot, B, BG, BM, E, OXF, MRDS 107314, 107315, 107337, 107360, 107376, 107422, 107428, 107429, 107436, 107498, 107538, 107546

Pyrenula parvinuclea (Meyen & Fw.) Aptroot

Uncommon, on *Barringtonia speciosa*, *Casuarina equisetifolia*, *Cocos nucifera*, *Guettarda speciosa* & *Scaevola sericea*.

B(S); DG; E; SI(F). Pantropical.

Herb. OXF, MRDS 107359, 107400, 107555

Pyxine cocoos (Sw.) Nyl.

Common, on *Argusia argentea*, *Cocos nucifera*, *Cordia subcordata*, *Guettarda speciosa* & *Hernandia sonora*.

B(M); B(N); B(S); DG; DG(I); E; N; SI(A); SI(B); SI(M); SI(S); SI(T). Cosmopolitan.

Herb. MRDS 107329, 107331, 107332, 107363, 107386, 107387, 107433, 107512, 107582

Pyxine consocians Vainio

Locally frequent, on *Cocos nucifera*.

DG; E; SI(DP); SI(B); SI(M). Pantropical.

Herb. OXF, MRDS 107372, 107511, 108798

Pyxine reticulata (Vainio) Vainio

Rare, on *Cocos nucifera*.

E. Pantropical.

Herb. MRDS 107402

Pyxine retirugella Nyl.

Rare, substratum unidentified.

DG; SI(M). Pantropical.

Rinodina sp.

Rare, on *Cocos nucifera*.

DG.

Siphula sp.

Rare, on thin poorly developed soil, covering several square metres.

SI(B).

Herb. Aptroot, MRDS 108110

Thelenella modesta (Nyl.) Nyl.

Rare, on lignum.

SI(M). Cosmopolitan.

Herb. E, MRDS 108229

Tomasellia eschweileri (Müll.Arg.) R.C.Harris

Rare, but overlooked, on *Cocos nucifera*. SI(B). Pantropical.

Herb. MRDS 107404

Verrucaria cf. maculiformis Krempelh.

Rare, on gravestone in abandoned plantation village.


DG. Cosmopolitan.

Geographical Affinities of the Flora

As can be seen from above, the lichen flora of Chagos mainly comprises pantropical species. The flora is characteristic of coastal tropical environments, and highly comparable to other small oceanic islands because of the absence of mountain species and species restricted to dense rainforest. The Chagos flora mainly consists of crustose lichens, except for the Physciaceae (syn. Pyxinaceae). As knowledge of tropical crustose lichens is still very incomplete, it is difficult to compare total species lists between different areas as the method of sampling and identification varies.

Nevertheless, a comparison can be made with some smaller islands which have been as exhaustively sampled by the authors as Chagos, namely the Maldives in the Indian Ocean (Aptroot 1991) and Laing Island in the Pacific Ocean (Aptroot *et al.* 1995). Less complete species lists are available for most smaller island groups in the Pacific Ocean (Elix & McCarthy 1998) and for Aldabra in the Indian Ocean (Seaward *et al.* 1996). For further comparisons, emphasis has been laid on some dominant elements in the Chagos flora, ie. Physciaceae and species growing on *Cocos*. Furthermore, comparisons could be made with neotropical sites, but no suitable species lists have been published in recent years. Therefore, a comparison is made with the epiphytic flora on *Cocos* in Belize (Barclay-Estrup 1992) and with unpublished 1998 records from Guadeloupe (Aptroot, unpublished data).

Relationships between ten different island and coastal tropical floras were tested by means



size on the number of lichen taxa (see also Table 1).

of the Sørensen Coefficient (K):

$$K = \frac{200 \times c}{a + b}$$

where a = number of species in one region, b = number of species in another region, and c = number of species common to both regions. As can be seen from Table 2, good correlations of lichen floras occur within the Indian Ocean, as would be expected, but poorer correlations existed within and between other geographical regions. Surprisingly, in the case of Chagos, its lichen flora showed an interesting ranking with other floras; in view of the predominating pantropical element, an overall more or less equal similarity between the floras was expected. Wide-ranging ranked correlations between Chagos and other floras are

in the sequence Maldives > Laing Island > Aldabra > Tuamotu > Pitcairn > N.Mariana & Belize > Guadeloupe > Cook. Although comparison of the Physciaceae of the ten floras (Table 2) show similarly high correlations between the Indian Ocean islands, the overall ranking is completely different. Comparisons with the Physciaceae floras of Madagascar (Aptroot 1988), Colombia (Aptroot 1989) and the Guianas (Aptroot 1987) have yet to be undertaken.

It is therefore concluded that although the lichen flora of Chagos is characteristic for an Indian Ocean island, it is dominated by pantropical species.

Acknowledgements

The senior author (MRDS) is particularly grateful to The Royal Society, British Ecological Society, The

Percy Sladen Memorial Fund, The Colin Reid Countryside Trust and the University of Bradford for funding which allowed him to participate in the 1996 Chagos Expedition; he also acknowledges with thanks the support of expedition members, particularly Prof.D.Bellamy, Dr C.R.C. Sheppard, Mr J.M.W. Topp, Cmdr N. Wells, Miss U. Gregory and Dr F. Stewart. Thanks are also due to Dr J. Egea (*Opegrapha*), Dr S. Ekman (*Bacidia*), Dr T.Lumbsch (*Lecanora*), Dr E. Serusiaux (*Fellhanera*) and Dr H.J.M. Sipman (foliicolous lichens) for identifying or confirming critical taxa as indicated, and to Mr S. Davidson and Dr A.D. Headley for their preparation of Figures 1 and 2 respectively.

References

- Aptroot,A. 1987.** Pyxinaceae (Lichens). *Flora of the Guianas*, series E, 1: 1-59.
- Aptroot,A. 1988.** Lichens of Madagascar: the Pyxinaceae (syn. Physciaceae). *Cryptogamie, Bryol.Lichenol.* 9: 141-147.
- Aptroot,A. 1989.** Studies on Colombian cryptogams. XL. The family Pyxinaceae (Lichenized Fungi). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen*, series C, 92: 269-280.
- Aptroot,A. 1991.** The lichen flora of The Maldives. *Lichenologist* 23: 57-60.
- Aptroot,A., Diederich,P., Serusiaux,E. & Sipman,H. 1995.** Lichens and lichenicolous fungi of Laing Island (Papua New Guinea). *Bibliotheca Lichenologica* 57: 19-48.
- Barclay-Estrup,P. 1992.** Lichens of *Cocos nucifera* on Caye Caulker, Belize. *Lichenologist* 24: 249-254.
- David,J.C. & Hawksworth,D.L. 1995.** Lichens of Mauritius I: some new species and records. *Bibliotheca Lichenologica* 57: 93-111.
- Eisenhauer,A., Heiss,G.A., Sheppard,C.R.C. & C.Dullo,W.-C. 1999.** Reef and island formation and Late Holocene sea-level changes in the Chagos islands. In: *Ecology of the Chagos Archipelago* (ed. C.R.C. Sheppard & M.R.D. Seaward): 21-33. Westbury Press/Linnean Society, London.
- Elix,J.A. & McCarthy,P.M. 1998.** Catalogue of the lichens of the smaller Pacific islands. *Bibliotheca Lichenologica* 70: 1-361.
- Hoffmann,L. 1989.** Algae of terrestrial habitats. *Botanical Reviews* 55: 77-105.
- Pocs,T. 1982.** Tropical forest bryophytes. In: *Bryophyte Ecology* (ed. A.J.E. Smith): 59-104. Chapman & Hall, London.
- Renvoize,S.A. 1975.** A floristic analysis of the western Indian Ocean coral islands. *Kew Bulletin* 30: 133-152.
- Richardson,D.H.S. 1981.** *The Biology of Mosses*. Blackwell, Oxford.
- Rogers,G.A. & Henriksson,E. 1976.** Associations between the blue-green algae *Anabaena variabilis* and *Nostoc muscorum* and the moss *Funaria hygrometrica* with reference to the colonization of Surtsey. *Acta Botanica Islandica* 4: 10-15.
- Seaward,M.R.D. 1999.** Cryptogamic flora of the Chagos Archipelago. In: *Ecology of the Chagos Archipelago* (ed. C.R.C. Sheppard & M.R.D. Seaward): 207-224. Westbury Press/Linnean Society, London.
- Seaward,M.R.D., Hambler,C. & Aptroot,A. 1996.** Bryophytes and lichens of Aldabra. *Tropical Bryology* 12: 29-33.
- Sheppard,C.R.C. & Seaward,M.R.D.,ed. 1999.** *Ecology of the Chagos Archipelago*. Westbury Press/Linnean Society, London.
- Sipman,H.J.M. & Harris,R.C. 1989.** Lichens. In: *Tropical Rain Forest Ecosystems* (ed. H. Lieth & M.J.A. Werger): 303-309. Elsevier, Amsterdam.
- Topp,J.M.W. & Sheppard,C.R.C. 1999.** Higher plants of the Chagos Archipelago. In: *Ecology of the Chagos Archipelago* (ed. C.R.C. Sheppard & M.R.D. Seaward): 225-240. Westbury Press/Linnean Society, London.
- Townsend,C.C. 1971.** List of Diego Garcia bryophyta. *Atoll Research Bulletin* 149: 161-162.